



COMMON PRE-BOARD EXAMINATION

CHEMISTRY-Code No. 043

Class-XII-(2025-26)



SET: 1




Time allowed: 3 Hrs.



Maximum Marks: 70

MARKING SCHEME

Q. No.	Section-A	Marks
1.	C) it shows positive deviation from Raoult's law	1
2	C) 10^{10}	1
3	(D) Cr^{3+}	1
4	B) Geometrical isomerism	1
5	D) P = $\text{CH}_3\text{-CH}_2\text{-CH}_2\text{-NO}_2$, Q = $\text{CH}_3\text{-CH}_2\text{-CH}_2\text{-NC}$	1
6	A) 	1
7	C) electrophilic substitution reaction	1
8	B) 2-methylpropene is formed	1
9	A) Sodium bicarbonate test	1
10	A) 	1
11	(D) $\text{CH}_3\text{CH}_2\text{NH}_2$	1
12	A) A-IV, B-III, C-I, D-II	1
13	B) Both A and R are true, and R is not the correct explanation of A.	1
14	A) Both A and R are true, and R is the correct explanation of A.	1
15	B) Both A and R are true, and R is not the correct explanation of A.	1
16	C) A is true but R is false.	1


Section-C		
Question No. 22 to 28 are short answer questions, carrying 3 marks each.		
22	$\text{KCl} \rightarrow \text{K}^+ + \text{Cl}^-$ $n = 2$ $i = 1 - \alpha + n \alpha$ $\Delta T_f = i K_f m$ $= (1 + 0.92) \times 1.86 \times \frac{0.5 \times 1000}{74.5 \times 100}$ $\Delta T_f = 0.24$ $\Delta T_f = T_f^0 - T_f^i$ $T_f^i = - 0.24 \text{ } ^\circ\text{C}$	 1/2 1/2 1/2 1/2 1/2 1/2
23	<p>(a) Yes, if the concentration of ZnSO₄ in the two half cell is different , the electrode potential will be different making the cell possible. (1)</p> <p>(b) $\Lambda^0_m (\text{MgCl}_2) = \lambda^0_m (\text{Mg}^{2+}) + 2 \lambda^0_m (\text{Cl}^-)$ $258.6 = 106 + 2 \lambda^0_m (\text{Cl}^-)$ $\lambda^0_m (\text{Cl}^-) = 76.3 \text{ Scm}^2\text{mol}^{-1}$</p> <p>(c) At anode : $\text{Zn}(\text{Hg}) + 2\text{OH}^- \rightarrow \text{ZnO} + \text{H}_2\text{O} + 2e^-$ At cathode: $\text{HgO} + \text{H}_2\text{O} + 2e^- \rightarrow \text{Hg} + 2\text{OH}^-$</p>	 1 1/2 1/2 1/2 1/2
24	<p>a) Cu^{2+} oxidizes iodide ion to iodine.</p> <p>b) Because of large number of unpaired electrons in their atoms they have stronger interatomic interaction or strong metallic bonding</p> <p>c) The chromates and dichromates are interconvertible in aqueous solution depending upon pH of the solution. Increasing the pH (in basic solution) of dichromate ions a colour change from orange to yellow is observed as dichromate ions change to chromate ions.</p>	 1 1 1
25	<p>a) $3\text{MnO}_4^{2-} + 4\text{H}^+ \rightarrow 2\text{MnO}_4^- + \text{MnO}_2 + 2\text{H}_2\text{O}$</p> <p>b) $\text{KMnO}_4 \xrightarrow{\Delta} \text{K}_2\text{MnO}_4 + \text{MnO}_2 + \text{O}_2$</p> <p>c) $5\text{Fe}^{2+} + \text{MnO}_4^- + 8\text{H}^+ \rightarrow 5\text{Fe}^{3+} + \text{Mn}^{2+} + 4\text{H}_2\text{O}$</p>	 1 1 1

26	$\begin{array}{c} \text{Cl} \\ \\ \text{CH}_3\text{-CH-CH}_2\text{Cl} \end{array}$ $\text{CH}_2\text{Cl-CH}_2\text{-CH}_2\text{Cl}$ $\text{CH}_3\text{-CH}_2\text{-CHCl}_2$ $\begin{array}{c} \text{Cl} \\ \\ \text{CH}_3\text{-C-CH}_3 \\ \\ \text{Cl} \end{array}$ <p>The following isomer will exhibit enantiomerism:</p> $\begin{array}{c} \text{Cl} \\ \\ \text{CH}_3\text{-CH-CH}_2\text{Cl} \end{array}$ <p>IUPAC name: 1,2-Dichloropropane.</p>	$\frac{1}{2} \times 6$
27	<p>(a)</p> $\text{H}_2\text{O} + \text{H}^+ \rightarrow \text{H}_3\text{O}^+$ $\text{>C=C<} + \text{H}-\overset{\text{H}}{\underset{\cdot\cdot}{\text{O}}^+}-\text{H} \rightleftharpoons \begin{array}{c} \text{H} \\ \\ \text{-C-} \end{array} \overset{+}{\text{C}}\text{<} + \text{H}_2\ddot{\text{O}}$ $\begin{array}{c} \text{H} \\ \\ \text{-C-} \end{array} \overset{+}{\text{C}}\text{<} + \text{H}_2\ddot{\text{O}} \rightleftharpoons \begin{array}{c} \text{H} \\ \\ \text{-C-} \end{array} \begin{array}{c} \text{H} \\ \\ \text{-C-} \end{array} \overset{+}{\text{O}}-\text{H}$ $\begin{array}{c} \text{H} \\ \\ \text{-C-} \end{array} \begin{array}{c} \text{H} \\ \\ \text{-C-} \end{array} \overset{+}{\text{O}}-\text{H} + \text{H}_2\ddot{\text{O}} \rightarrow \begin{array}{c} \text{H} \\ \\ \text{-C-} \end{array} \begin{array}{c} \text{:OH} \\ \\ \text{-C-} \end{array} + \text{H}_3\text{O}^+$ <p>(b)</p> 	$\frac{1}{2}$ 1 $\frac{1}{2}$ $\frac{1}{2} + \frac{1}{2}$
28	<p>a)</p> $\text{>C=O} \xrightarrow[-\text{H}_2\text{O}]{\text{NH}_2\text{NH}_2} \text{>C=NNH}_2 \xrightarrow[\text{heat}]{\text{KOH/ethylene glycol}} \text{>CH}_2 + \text{N}_2$ <p>b)</p> $\text{R-CH}_2\text{-COOH} \xrightarrow[\text{(ii) H}_2\text{O}]{\text{(i) X}_2/\text{Red phosphorus}} \begin{array}{c} \text{R-CH-COOH} \\ \\ \text{X} \\ \text{X = Cl, Br} \end{array}$	1 1

	<p>c)</p>  <p>Formaldehyde + Conc. KOH $\xrightarrow{\Delta}$ Methanol + Potassium formate</p> <p>d) $\text{CH}_3\text{CN} + \text{SnCl}_2 + \text{HCl} \rightarrow \text{CH}_3\text{-CH=NH} \xrightarrow{\text{H}_3\text{O}^+} \text{CH}_3\text{CHO}$</p> <p>(Any three)</p>	<p>1</p> <p>1</p>
<p>Section D</p> <p>Question No. 29 & 30 are case-based/data -based questions carrying 4 marks each.</p>		
<p>29</p>	<p>(i) Secondary valency = 4</p> <p>(ii) cis form of $[\text{PtCl}_2(\text{en})_2]^{2+}$ shows optical isomerism</p>  <p>(iii) A (1) $\text{Fe}_4[\text{Fe}(\text{CN})_6]_3$ (2) $t_2g^3 e_g^2$</p> <p>OR</p> <p>(iii) B. dsp^2, diamagnetic</p>	<p>1</p> <p>1</p> <p>1</p> <p>1 + 1</p>
<p>30</p>	<p>(i) The amounts of different substances liberated by the same quantity of electricity passing through the electrolytic solution are proportional to their chemical equivalent weights</p> <p>(ii) When an external potential greater than cell potential is applied.</p> <p>(iii) A.</p> $m = z I t$ $2.8 \text{ g} = \frac{56 \times 2 \times t}{2 \times 96500}$ $t = 4825 \text{ s}$ <p style="text-align: center;">OR</p> <p>(iii) B</p> $E_{\text{cell}} = E^{\circ}_{\text{cell}} - \frac{0.059}{n} \log Q_c$ $= E^{\circ}_{\text{cell}} - \frac{0.059}{2} \log \frac{10^{-3}}{10^{-2}}$ $= 2.71 + 0.0295$ $E_{\text{cell}} = 2.7395 \text{ V}$	<p>1</p> <p>1</p> <p>$\frac{1}{2}$</p> <p>1</p> <p>$\frac{1}{2}$</p> <p>$\frac{1}{2}$</p> <p>$\frac{1}{2}$</p> <p>$\frac{1}{2}$</p>


	<p>d)</p> $ \begin{array}{ccc} \text{CHO} & & \text{COOH} \\ & & \\ (\text{CHOH})_4 & \xrightarrow{\text{Br}_2 \text{ water}} & (\text{CHOH})_4 \\ & & \\ \text{CH}_2\text{OH} & & \text{CH}_2\text{OH} \\ & & \text{Gluconic acid} \end{array} $ $ \begin{array}{ccc} \text{CHO} & & \text{COOH} \\ & & \\ (\text{CHOH})_4 & \xrightarrow{\text{Oxidation}} & (\text{CHOH})_4 \\ & & \\ \text{CH}_2\text{OH} & & \text{COOH} \\ & & \text{Saccharic acid} \end{array} $ <p style="text-align: center;">OR</p> <p>B. (a) This indicates that the aldehyde group in glucose is not free. (b)</p> <p style="text-align: center;">α-D-(+)-Glucopyranose</p> <p>(c) 'D' gives the configuration i.e. the – OH gp at carbon 5 is on the right hand side. (+) indicates that the isomer is dextro rotatory. (d) Vitamin B₂ , Milk , Egg white , liver, kidney (any two)</p>	<p>1</p> <p>1</p> <p>1</p> <p>1</p> <p>1/2</p> <p>1/2</p> <p>1 + 1</p>
33	<p>A a)</p> <p>(1)</p> <p>(2)</p> <p>(3)</p> <p>b) Aniline undergoes resonance and as a result the electrons on the N-atom are less available for donation. c) $(\text{CH}_3)_3\text{N} < \text{CH}_3\text{NH}_2 < (\text{CH}_3)_2\text{NH}$</p>	<p>1</p> <p>1</p> <p>1</p> <p>1</p> <p>1</p> <p>1</p>

B. a) (i)




1 + 1

(ii)




1

b)



1

(ii)



1